

J-CARAF 全8回調査の概要

(グレイの部分は調査項目に含まれていません。赤字は2020.2.20に訂正)

Table with columns for survey order (1-8), target period (2011-2017), and percentages. Rows include categories like '概要' (Summary), '併存症など' (Comorbidities), '心エコー' (Echocardiography), '抗不整脈薬治療の経験' (Anti-arrhythmic drug experience), '画像検査' (Imaging), '術時間' (Procedure time), '使用機器' (Equipment), '肺静脈隔離の方法' (PVI method), and 'PVI以外の治療' (Non-PVI treatments).

	Amiodarone	50	71	76	56	85	122	122	119	5.4%	5.9%	6.1%	5.3%	4.8%	5.8%	4.9%	4.9%
	その他	14	20	15	3	12	18	20	40	1.5%	1.7%	1.2%	0.3%	0.7%	0.9%	0.8%	1.6%
	抗不整脈薬の併用	46	51	60	17	46	65	48	3	4.9%	4.2%	4.8%	1.6%	2.6%	3.1%	1.9%	0.1%
	Verapamil	64	68	73	49	95	69	103	55	6.9%	5.6%	5.8%	4.7%	5.3%	3.3%	4.2%	2.3%
	β blockade	253	375	350	309	601	766	938	964	27.1%	31.1%	28.0%	29.5%	33.8%	36.4%	37.9%	39.5%
	ACEI	51	68	53	40	68	113	154	160	5.5%	5.6%	4.2%	3.8%	3.8%	5.4%	6.2%	6.6%
	ARB	233	284	264	213	401	389	573	534	25.0%	23.6%	21.1%	20.3%	22.6%	18.5%	23.1%	21.9%
	Statin	153	208	139	156	304	318	494	429	16.4%	17.3%	11.1%	14.9%	17.1%	15.1%	20.0%	17.6%